

## How to Manage Pests

### UC Pest Management Guidelines

| [More pests](#) | [More crops](#) | [About guidelines](#) |

## Tomato Verticillium Wilt

**Pathogen:** *Verticillium dahliae*

(Reviewed 1/07, updated 1/07)

In this Guideline:

- [Symptoms](#)
- [Publication](#)
- [Comments on the disease](#)
- [Glossary](#)
- [Management](#)



### SYMPTOMS AND SIGNS

Older leaves on tomato plants infected with *Verticillium* appear as yellow, V-shaped areas that narrow from the margin. The leaf progressively turns from [yellow to brown](#) and eventually dies. Older and lower leaves are the most affected. Sun-related fruit damage is increased because of the loss of foliage. A light tan [discoloration](#) develops in the vascular tissue, especially near the base of the plant. The discoloration extends a short distance up the plant and may occur in patches. Symptoms are most noticeable during later stages of plant development when fruit begin to size.

### COMMENTS ON THE DISEASE

The fungus survives as microsclerotia in the soil. Once established in a field, it persists indefinitely and can cause disease whenever a susceptible host is planted. A large number of crops and weeds serve as hosts. The disease is favored by cool soil and air temperatures. Verticillium wilt is difficult to distinguish from [Fusarium wilt](#) and positive identification may require cultivating the fungus in a laboratory. Verticillium wilt seldom kills tomato plants but reduces their vigor and yield.

### MANAGEMENT

Use resistant cultivars effective against Race 1. No source of resistance to Race 2 is commercially available. Sanitation, especially washing equipment to prevent movement of infested soil, may help to slow spread of the Race 2 strain of the pathogen. Rotation to nonsusceptible crops, such as small grains and corn, helps reduce inoculum.

### PUBLICATION



*UC IPM Pest Management Guidelines: Tomato*

UC ANR Publication 3470

Diseases

R. M. Davis, Plant Pathology, UC Davis

G. Miyao, UC Cooperative Extension, Solano/Yolo counties

K. Subbarao, USDA Agricultural Research Station, Salinas

J. J. Stapleton, UC IPM Program, Kearney Agricultural Center, Parlier

Acknowledgments for contributions to the disease section:

B. W. Falk, Plant Pathology, UC Davis

Statewide IPM Program, Agriculture and Natural Resources, University of California

All contents copyright © 2012 The Regents of the University of California. All rights reserved.

For noncommercial purposes only, any Web site may link directly to this page. FOR ALL OTHER USES or more information, read [Legal Notices](#). Unfortunately, we cannot provide individual solutions to specific pest problems. See our [Home page](#), or in the U.S., contact your [local Cooperative Extension office](#) for assistance. /PMG/r783100911.html revised: March 16, 2012. [Contact webmaster](#).